


We claim:

1. A fungicidal mixture, comprising as active components

(1) a benzamidoxime derivative of the formula I




where the substituent and the index may have the following meanings:

R is hydrogen, halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy or C<sub>1</sub>-C<sub>4</sub>-haloalkoxy,

n is 1, 2 or 3,

and anazole derivative or a salt or adduct thereof, selected from the group consisting of


(2) bromuconazole of the formula II



and


(3) cyproconazole of the formula III

24




and

10 (4) difenoconazole of the formula IV




and

20 (5) diniconazole of the formula V



and

30 (6) epoxiconazole of the formula VI




and

45 (7) fenbuconazole of the formula VII

25

5




(VII)

and

10

(8) fluquinconazole of the formula VIII

15




(VIII)

20

and

(9) flusilazole of the formula IX

25




(IX)

30

and

(10) hexaconazole of the formula X

35



(X)


40

and

(11) metconazole of the formula XI

45


26



(XI)

and


(12) prochloraz of the formula XII



(XII)

and


(13) propiconazole of the formula XIII



(XIII)

and

(14) tebuconazole of the formula XIV




(XIV)

and

(15) tetraconazole of the formula XV


45

27




and

10 (16) triflumizole of the formula XVI




and

20 (17) flutriafol of the formula XVII



and


30 (18) myclobutanil of the formula XVIII



and

45 (19) penconazole of the formula XIX


28



(XIX)

and


(20) simeconazole of the formula XX



(XX)

and

(21) ipconazole of the formula XXI




(XXI)

and

(22) triticonazole of the formula XXII

29

5




(XXII)

10

and

(23) prothioconazole of the formula XXIII

15



(XXIII)

20

25 in a synergistically effective amount.

2. A fungicidal mixture as claimed in claim 1 where in formula I the radical R is hydrogen.
- 30 3. A fungicidal mixture as claimed in claim 1, wherein the weight ratio of the benzamidoxime derivative of the formula I to the triazole of the formulae II to XXIII in question is from 20 : 1 to 1 : 20.
- 35 4. A method for controlling harmful fungi, which comprises treating the harmful fungi, their habitat or the plants, seeds, soils, areas, materials or spaces to be kept free from them with a fungicidal mixture as claimed in claim 1.
- 40 5. A method as claimed in claim 4, wherein the compound of the formula I as set forth in claim 1 and at least one compound of the formulae II to XXIII as set forth in claim 1 are applied simultaneously, i.e. together or separately, or successively.

45

30

6. A method as claimed in claim 4 or 5, wherein the fungicidal mixture or the compound of the formula I and at least one compound of the formulae II to XXIII as set forth in claim 1 are applied in an amount of from 0.01 to 8 kg/ha.

5

7. A fungicidal composition, comprising the fungicidal mixture as claimed in claim 1 and a solid or liquid carrier.

10

15

20

25

30

35

40

45